

International Journal of Modern Physics A
 © World Scientific Publishing Company

B MESON LIGHT-CONE DISTRIBUTION AMPLITUDES AND HEAVY-QUARK SYMMETRY*

HIROYUKI KAWAMURA

*Deutsches Elektronen-Synchrotron, DESY
 Platanenallee 6, D 15738 Zeuthen, GERMANY*

JIRO KODAIRA and CONG-FENG QIAO[†]

*Dept. of Physics, Hiroshima University
 Higashi-Hiroshima 739-8526, JAPAN*

KAZUHIRO TANAKA

*Dept. of Physics, Juntendo University
 Inba-gun, Chiba 270-1695, JAPAN*

We present a systematic study of the B meson light-cone distribution amplitudes which are relevant for the QCD factorization approach for the exclusive B meson decays. We construct representations for the quark-antiquark distribution amplitudes in terms of independent dynamical degrees of freedom, which exactly satisfy the QCD equations of motion and constraints from heavy-quark symmetry.

Recently systematic methods based on the QCD factorization have been developed for the exclusive B meson decays into light mesons^{1,2,3,4} (for other approaches see⁵). Essential ingredients in this approach are the light-cone distribution amplitudes for the participating mesons, which constitute nonperturbative long-distance contribution to the factorized amplitudes. For the light mesons (π , K , ρ , K^* , etc.) appearing in the final state, systematic model-independent study of the light-cone distributions exists for both leading and higher twists.⁶ On the other hand, the light-cone distribution amplitudes for the B meson are not well-known at present and provide a major source of uncertainty in the calculations of the decay rates. In this work,⁷ we demonstrate that heavy-quark symmetry and constraints from the equations of motion determine a unique analytic solution for the B meson light-cone distribution amplitudes within the two-particle Fock states. We also derive the exact integral representations for the effects of higher Fock states with additional gluons.

In the heavy-quark limit, the B meson matrix elements obey the heavy-quark symmetry, and is described by the heavy-quark effective theory (HQET).⁸ Following

*Talk presented by K. Tanaka.

[†]JSPS Research Fellow.

2 *H. Kawamura, J. Kodaira, C.F. Qiao and K. Tanaka*

Refs.^{9,2}, we introduce the quark-antiquark light-cone distribution amplitudes $\tilde{\phi}_{\pm}(t)$ of the B meson in terms of matrix element in the HQET:

$$\langle 0 | \bar{q}(z) \Gamma h_v(0) | \bar{B}(p) \rangle = -\frac{if_B M}{2} \text{Tr} \left[\gamma_5 \Gamma \frac{1 + \not{v}}{2} \left\{ \tilde{\phi}_+(t) - \not{z} \frac{\tilde{\phi}_+(t) - \tilde{\phi}_-(t)}{2t} \right\} \right] . \quad (1)$$

where z_μ is a light-like vector ($z^2 = 0$), $v^2 = 1$, $t = v \cdot z$, and $p^\mu = Mv^\mu$ is the 4-momentum of the B meson with mass M . $h_v(x)$ denotes the effective b -quark field,⁸ $b(x) \approx \exp(-im_b v \cdot x) h_v(x)$, and is subject to the on-shell constraint, $\not{v} h_v = h_v$. Γ is a generic Dirac matrix and, here and in the following, the path-ordered gauge factors are implied in between the constituent fields. f_B is the decay constant defined as usually as $\langle 0 | \bar{q}(0) \gamma^\mu \gamma_5 h_v(0) | \bar{B}(p) \rangle = if_B M v^\mu$ so that $\tilde{\phi}_{\pm}(t=0) = 1$. $\tilde{\phi}_+$ is of leading-twist, whereas $\tilde{\phi}_-$ has subleading twist.⁹

It is well-known that the QCD equations of motion impose a set of relations between distribution amplitudes for the light-mesons.⁶ The corresponding relations can be derived from the identities between the nonlocal operators:

$$\frac{\partial}{\partial x^\mu} \bar{q}(x) \gamma^\mu \Gamma h_v(0) = i \int_0^1 du u \bar{q}(x) g G_{\mu\nu}(ux) x^\nu \gamma^\mu \Gamma h_v(0) , \quad (2)$$

$$v^\mu \frac{\partial}{\partial x^\mu} \bar{q}(x) \Gamma h_v(0) = i \int_0^1 du (u-1) \bar{q}(x) g G_{\mu\nu}(ux) v^\mu x^\nu \Gamma h_v(0) + v^\mu \frac{\partial}{\partial y_\mu} \bar{q}(x+y) \Gamma h_v(y) \Big|_{y \rightarrow 0} , \quad (3)$$

where $G_{\mu\nu}$ is the gluon field strength tensor, and we have used the equations of motion $\not{p} q = 0$ and $v \cdot D h_v = 0$ with $D_\mu = \partial_\mu - ig A_\mu$ the covariant derivative. We take the vacuum-to-meson matrix element of these identities and go over to the light-cone limit $x_\mu \rightarrow z_\mu$. The terms in the LHS of eqs. (2) and (3) yield $\tilde{\phi}_+(t)$, $\tilde{\phi}_-(t)$ defined above and their derivatives: $d\tilde{\phi}_{\pm}(t)/dt$ and $\partial\tilde{\phi}_{\pm}(t, x^2)/\partial x^2|_{x^2 \rightarrow 0}$, where, via $\tilde{\phi}_{\pm}(t) \rightarrow \tilde{\phi}_{\pm}(t, x^2)$, we extend the definitions in eq.(1) to the case $z \rightarrow x$ ($x^2 \neq 0$). The last term of (3), the derivative over total translation, yields contribution with

$$\bar{\Lambda} = M - m_b = \frac{iv \cdot \partial \langle 0 | \bar{q} \Gamma h_v | \bar{B}(p) \rangle}{\langle 0 | \bar{q} \Gamma h_v | \bar{B}(p) \rangle} . \quad (4)$$

This is the usual “effective mass” of meson states in the HQET.⁸

The terms given by an integral of quark-antiquark-gluon operator are expressed by the three-particle distribution amplitudes corresponding to the higher-Fock components of the meson wave function. Through the Lorentz decomposition of the three-particle light-cone matrix element, we define the three functions $\tilde{\Psi}_V(t, u)$, $\tilde{\Psi}_A(t, u)$, $\tilde{X}_A(t, u)$ as the independent three-particle distributions:⁷

$$\begin{aligned} & \langle 0 | \bar{q}(z) g G_{\mu\nu}(uz) z^\nu \Gamma h_v(0) | \bar{B}(p) \rangle \\ &= \frac{1}{2} f_B M \text{Tr} \left[\gamma_5 \Gamma \frac{1 + \not{v}}{2} \left\{ (v_\mu \not{z} - t \gamma_\mu) \left(\tilde{\Psi}_A(t, u) - \tilde{\Psi}_V(t, u) \right) \right. \right. \\ & \quad \left. \left. - i \sigma_{\mu\nu} z^\nu \tilde{\Psi}_V(t, u) + \frac{z_\mu}{t} (\not{z} - t) \tilde{X}_A(t, u) \right\} \right] . \quad (5) \end{aligned}$$

From the two identities (2) and (3), we eventually obtain the four independent differential equations for the distribution amplitudes,⁷ which are exact in QCD in the heavy-quark limit. This system of four differential equations can be organized into the two sets, so that the first set of two equations does not involve the derivatives with respect to the transverse separation, $\partial\tilde{\phi}_{\pm}(t, x^2)/\partial x^2|_{x^2 \rightarrow 0}$, while the second set of two equations involves them; the second set is uninteresting for our purpose. The first set of equations is given by⁷

$$\omega \frac{d\phi_{-}(\omega)}{d\omega} + \phi_{+}(\omega) = I(\omega) , \quad (6)$$

$$(\omega - 2\bar{\Lambda}) \phi_{+}(\omega) + \omega \phi_{-}(\omega) = J(\omega) , \quad (7)$$

where we have introduced the momentum-space distribution amplitudes $\phi_{\pm}(\omega)$ as $\tilde{\phi}_{\pm}(t) = \int d\omega e^{-i\omega t} \phi_{\pm}(\omega)$ with ωv^{+} the light-cone projection of the light-antiquark momentum in the B meson. $I(\omega)$ and $J(\omega)$ of eqs. (6) and (7) denote the “source” terms due to three-particle amplitudes as⁷

$$I(\omega) = 2 \frac{d}{d\omega} \int_0^{\omega} d\rho \int_{\omega-\rho}^{\infty} \frac{d\xi}{\xi} \frac{\partial}{\partial \xi} [\Psi_A(\rho, \xi) - \Psi_V(\rho, \xi)] , \quad (8)$$

and similarly for $J(\omega)$. Here the three-particle amplitudes in the momentum space are defined as $\tilde{F}(t, u) = \int d\omega d\xi e^{-i(\omega+\xi)u} F(\omega, \xi)$ with $F = \{\Psi_V, \Psi_A, X_A\}$.

A system of equations (6), (7) can be solved for $\phi_{+}(\omega)$ and $\phi_{-}(\omega)$ with boundary conditions $\phi_{\pm}(\omega) = 0$ for $\omega < 0$ or $\omega \rightarrow \infty$, and with normalization condition $\int_0^{\infty} d\omega \phi_{\pm}(\omega) = \tilde{\phi}_{\pm}(0) = 1$. The solution can be decomposed into two pieces as

$$\phi_{\pm}(\omega) = \phi_{\pm}^{(WW)}(\omega) + \phi_{\pm}^{(g)}(\omega) , \quad (9)$$

where $\phi_{\pm}^{(WW)}(\omega)$ are the solution of eqs. (6) and (7) with $I(\omega) = J(\omega) = 0$, which corresponds to $\Psi_V = \Psi_A = X_A = 0$ (“Wandzura-Wilczek approximation”). $\phi_{\pm}^{(g)}(\omega)$ denote the pieces induced by the source terms.

Eq.(6) alone, with $I(\omega) = 0$, is equivalent to a usual Wandzura-Wilczek type relation derived in Ref.²: $\phi_{-}^{(WW)}(\omega) = \int_{\omega}^{\infty} d\rho \phi_{+}^{(WW)}(\rho)/\rho$. Combining eqs.(6) and (7), we are able to obtain the analytic solution explicitly as ($\omega \geq 0$)

$$\phi_{+}^{(WW)}(\omega) = \frac{\omega}{2\bar{\Lambda}^2} \theta(2\bar{\Lambda} - \omega) , \quad \phi_{-}^{(WW)}(\omega) = \frac{2\bar{\Lambda} - \omega}{2\bar{\Lambda}^2} \theta(2\bar{\Lambda} - \omega) . \quad (10)$$

The solution for $\phi_{\pm}^{(g)}$ can be obtained straightforwardly, and reads ($\omega \geq 0$):

$$\phi_{+}^{(g)}(\omega) = \frac{\omega}{2\bar{\Lambda}} \Phi(\omega) , \quad \phi_{-}^{(g)}(\omega) = \frac{2\bar{\Lambda} - \omega}{2\bar{\Lambda}} \Phi(\omega) + \frac{J(\omega)}{\omega} , \quad (11)$$

$$\begin{aligned} \Phi(\omega) = & \theta(2\bar{\Lambda} - \omega) \left\{ \int_0^{\omega} d\rho \frac{K(\rho)}{2\bar{\Lambda} - \rho} - \frac{J(0)}{2\bar{\Lambda}} \right\} - \theta(\omega - 2\bar{\Lambda}) \int_{\omega}^{\infty} d\rho \frac{K(\rho)}{2\bar{\Lambda} - \rho} \\ & - \int_{\omega}^{\infty} d\rho \left(\frac{K(\rho)}{\rho} + \frac{J(\rho)}{\rho^2} \right) , \end{aligned} \quad (12)$$

with $K(\rho) = I(\rho) + [1/(2\bar{\Lambda}) - d/d\rho] J(\rho)$. The solution (9) with eqs.(10)-(12) is exact and presents our principal result.

4 H. Kawamura, J. Kodaira, C.F. Qiao and K. Tanaka

It is also straightforward to derive the Mellin moments $\langle \omega^n \rangle_{\pm} \equiv \int d\omega \omega^n \phi_{\pm}(\omega)$ ($n = 0, 1, 2, \dots$) of our solution. Because the analytic expression for general moment n is somewhat complicated,⁷ we present some examples for a few low moments: $\langle \omega \rangle_+ = 4\bar{\Lambda}/3$, $\langle \omega \rangle_- = 2\bar{\Lambda}/3$, and

$$\langle \omega^2 \rangle_+ = 2\bar{\Lambda}^2 + \frac{2}{3}\lambda_E^2 + \frac{1}{3}\lambda_H^2, \quad \langle \omega^2 \rangle_- = \frac{2}{3}\bar{\Lambda}^2 + \frac{1}{3}\lambda_H^2, \quad (13)$$

where λ_E and λ_H are due to $\phi_{\pm}^{(g)}$, and are related to the chromoelectric and chromomagnetic fields in the B meson rest frame as $\langle 0 | \bar{q} g \mathbf{E} \cdot \boldsymbol{\alpha} \gamma_5 h_v | \bar{B}(\mathbf{p} = 0) \rangle = f_B M \lambda_E^2$, $\langle 0 | \bar{q} g \mathbf{H} \cdot \boldsymbol{\sigma} \gamma_5 h_v | \bar{B}(\mathbf{p} = 0) \rangle = i f_B M \lambda_H^2$. These results for $n = 1, 2$ coincide with the relations obtained by Grozin and Neubert,⁹ who have derived their relations by analyzing matrix elements of *local* operators corresponding to these moments. Our solution (9) from nonlocal operators gives generalization of theirs to $n \geq 3$.

Our results reveal that the leading-twist distribution amplitude ϕ_+ as well as the higher-twist ϕ_- contains the three-particle contributions, which is in contrast with the case of the light mesons.⁶ We note that there exists an estimate of λ_E and λ_H by QCD sum rules,⁹ but any estimate of the higher moments is not known. Further investigations are required to clarify the effects of multi-particle states. In this connection, we note that the shape of our Wandzura-Wilczek contributions (10), which are determined uniquely in analytic form in terms of $\bar{\Lambda}$, is rather different from various “model” distribution amplitudes that have been used in the existing literature (see Ref.⁹ and references therein).

To conclude, our solution provides the powerful framework for building up the B meson light-cone distribution amplitudes and their phenomenological applications, because the solution is exact and satisfies all relevant QCD constraints.

We thank the organizers and all others who helped make the symposium successful. We thank T. Onogi for fruitful discussions. The work of J.K. was supported in part by the Monbu-kagaku-sho Grant-in-Aid for Scientific Research No.C-13640289. The work of C-F.Q. was supported by the Grant-in-Aid of JSPS committee.

References

1. M. Beneke et al., *Nucl. Phys.* **B591**, 313 (2000).
2. M. Beneke and T. Feldmann, *Nucl. Phys.* **B592**, 3 (2001).
3. M. Beneke, T. Feldmann and D. Seidel, *Nucl. Phys.* **B612**, 25 (2001).
S. W. Bosch and G. Buchalla, hep-ph/0106081.
4. C. W. Bauer, D. Pirjol and I. W. Stewart, *Phys. Rev. Lett.* **87**, 201806 (2001).
5. P. Ball and V. M. Braun, *Phys. Rev.* **D58**, 094016 (1998).
Y. Y. Keum, H. Li and A. I. Sanda, *Phys. Rev.* **D63**, 054008 (2001).
6. P. Ball, V. M. Braun, Y. Koike and K. Tanaka, *Nucl. Phys.* **B529**, 323 (1998).
P. Ball, *JHEP* **9901**, 010 (1999), and references therein.
7. H. Kawamura, J. Kodaira, C.F. Qiao and K. Tanaka, *Phys. Lett.* **B523**, 111 (2001).
8. M. Neubert, *Phys. Rept.* **245**, 259 (1994), and references therein.
9. A. G. Grozin and M. Neubert, *Phys. Rev.* **D55**, 272 (1997).